



For more information about the Organization of Teratology Information Specialists or to find a service in your area, call (866) 626-6847 or visit us online at: www.OTISpregnancy.org.

Caffeine and Pregnancy

This sheet talks about the risks that exposure to caffeine can have during pregnancy. With each pregnancy, all women have a 3% to 5% chance of having a baby with a birth defect. This information should not take the place of medical care and advice from your health care provider.

What is caffeine?

Caffeine is a stimulant found in many foods and beverages. It is also found in prescription and over the counter medications. Caffeine is a naturally occurring substance found in the leaves, seeds, and fruits of more than 60 plants.

How much caffeine is in common foods and drinks?

A few examples of caffeinated items include: 8oz. cup of coffee (65-120mg of caffeine), brewed tea (20-90mg), 12oz. Coke® (45mg), 1 cup of Haagen-Dazs® coffee ice cream (58mg), Hershey® chocolate bar (10mg), 8oz. hot chocolate (5mg), and 2 tablets of Excedrin® (130mg). Some herbals such as guarana also contain caffeine.

What exactly does caffeine do to my body?

Caffeine's main effect is increased alertness. Within 1 hour of coffee consumption, caffeine usually reaches its peak level in the bloodstream and remains there for 4-6 hours. Caffeine also stimulates the release of acid in the stomach, sometimes resulting in an upset stomach. Caffeine is also a diuretic, which means it helps eliminate fluids from the body and can result in water and calcium loss.

I am trying to become pregnant. Is it true that drinking caffeinated beverages will lower my chance to become pregnant?

Results from studies have been mixed. Some studies have suggested that high levels of caffeine (>300mg/day) may make it harder to conceive, but these findings are not proven. Low to moderate caffeine consumption (<300mg/day) does not seem to reduce a woman's chance of becoming pregnant.

I am pregnant. Is it safe for me to continue drinking coffee and soda?

Most experts agree that moderation and common sense are the keys for consuming caffeinated items during pregnancy. "Moderate" caffeine consumption is approximately 300mg/day, which is similar to 3 cups of coffee. It is also important for pregnant women to drink enough water, milk and juice. These fluids should not be replaced with caffeinated beverages.

Does caffeine cause or contribute to miscarriage?

Researchers have had difficulty determining whether there is a relationship between caffeine and miscarriage since miscarriage is very common. Recent reports suggest that low to moderate consumption of caffeine does not increase the risk for miscarriage. A few studies have shown that

there may be an increased risk for miscarriage or fetal death with high caffeine consumption (>300 mg/day), particularly in combination with smoking or alcohol, or with very high levels of caffeine consumption (>800 mg/day).

Will drinking caffeinated beverages during my pregnancy cause birth defects in my baby?

No. In humans, even large amounts of caffeine have not been shown to cause an increased chance for birth defects.

I drink 5-6 cups of coffee a day and I am 7 months pregnant. Will this amount of caffeine affect my baby?

Caffeine does cross the placenta; therefore large amounts of caffeine could affect babies in the same way as it does adults. Some reports have stated that children born to mothers who consumed >500mg/day were more likely to have faster heart rates, tremors, increased breathing rate, and spend more time awake in the days following birth.

Is it a problem if the baby's father consumed a lot of caffeine when I became pregnant?

Very little information is available on the effects of caffeine on sperm. One report stated that caffeine seems to increase the motility (movement) of sperm. Currently there are no other known effects on pregnancy.

Can I drink caffeinated beverages while I breastfeed?

Caffeine passes into breast milk, but there is very little information on how it might affect the baby. Caffeine consumption should be limited while breastfeeding. Women should remain well hydrated with water, juice and milk while breastfeeding.

December 2006.

Copyright by OTIS.

Reproduced by permission.



References:

Bech BH, et al. 2005. Coffee and fetal death: a cohort study with prospective data. *Am J Epidemiol* 162(10):983-990.

Christian M and Brent R. 2001. Teratogen Update: Evaluation of the Reproductive and Developmental Risks of Caffeine. *Teratology* 64: 51-78.

Cnattingius S, et al. 2000. Caffeine Intake and the Risk of First-Trimester Spontaneous Abortion. *N Engl J Med* 343:1839-1845.

Eskenazi B 1999. Caffeine—Filtering the Facts. *N Engl J Med* 341:1688-1689.

Hatch E and Bracken M. 1993. Association of Delayed Conception with Caffeine Consumption. *Am J Epidemiol* 138:1082-1091.

Genetic Drift. 2000. Caffeine in Pregnancy: Is there an Increased Risk for Miscarriage? Vol 11. www.mostgene.org.

International Food Information Council Foundation. 1998. Everything You Need to Know About... Caffeine. <http://ificinfo.health.org>.

Klebanoff M, et al. 1999. Maternal Serum Paraxanthine, a Caffeine Metabolite, and the Risk of Spontaneous Abortion. *New Engl J Med* 341:1639-1644.

Larroque B, et al. 1993. Effects on Birthweight of Alcohol and Caffeine Consumption During Pregnancy. *Am J Epidemiol* 137:941-950.

Matijasevich A, et al. 2006. Maternal caffeine consumption and fetal death: a case-control study in Uruguay. *Paediatr Perinat Epidemiol* 20(2):100-109.

Mills JL, et al. 1993. Moderate Caffeine Use and the Risk of Spontaneous Abortion and Intrauterine Growth. *JAMA* 269:593-597.

Shu XO, et al. 1995. Maternal Smoking, Alcohol Drinking, Caffeine Consumption and Fetal Growth: Results from a Prospective Study. *Epidemiology* 6:115-120.

If you have questions about the information on this fact sheet or other exposures during pregnancy, call OTIS at 1-866-626-6847.